

Charged Particles in Electromagnetic Field in Classical Case

Jyotiranjana Mohanty¹ Swagatika Dash²

¹Asst.Professor, PHYSICS, GIFT College, BHUBANESWAR.

²Asst.Professor, PHYSICS, GEC, BBSR, Odisha.

ABSTRACT- The study of the physical system consisting of charged particles in electromagnetic field constitute a major part of the whole of physics. Herestarting with the general laws of classical electrodynamics in the covariant form and then consider special cases of uniform and non-uniform electromagnetic fields with examples to find the trajectories in exact form or in an approximation.

I. INTRODUCTION

Here we consider the motion of a charged particle in various electromagnetic fields in the absence of a medium. We follow the treatments of Landau-Lifshitz and Jackson here.

The equation of motion of a charge in an electromagnetic field can be written as

$$\frac{d\varepsilon_{kin}}{dt} = eE.v \quad \dots\dots\dots 1.1$$

$$\frac{dp}{dt} = eE + \frac{e}{c}V * H$$

The expression on the right of equation (1.1) is called **Lorentz Force**.

The work done on the charged particle by the electric field is given by

The magnetic field does no work on a charge moving in it because the force which the magnetic field exerts is always perpendicular to the velocity of the charge

$$\varepsilon = \frac{mc^2}{\sqrt{1 - \frac{v^2}{c^2}}} + e\phi \quad \dots\dots\dots 1.2$$

Hence the energy of a charged particle in a constant time independent electromagnetic field can be written as

The presence of the field adds to the energy of the particle i.e. $e\phi$ the potential energy of the charge in the field. Energy depends only scalar but not on the vector potential. This means that the magnetic field

$$m \frac{d^\alpha X^\alpha}{d\tau^2} = \frac{q}{c} (\partial^\alpha A^\beta - \partial^\beta A^\alpha) \frac{dX_\beta}{d\tau} \quad \dots\dots\dots 1.3$$

does not affect the energy of the charge, only electric field can change the energy.

The covariant equation of motion of a charged particle can be written as :

Where τ is the proper time

1.1 MOTION IN CONSTANT ELECTRIC AND MAGNETIC FIELDS

Consider the motion of a charged particle 'e' moving in a combination of electric and

$$m\dot{v} = eE + \frac{e}{c}V * H \quad \dots\dots\dots 1.4$$

magnetic fields E and H, both uniform and constant. For this the equation of motion along the direction of H as z – axis will be

$$m\ddot{x} = \frac{e}{c} \dot{y}H$$

or
$$m\ddot{y} = eEy \frac{e}{c} \dot{x}H \quad \dots\dots\dots 1.12$$

$$m\ddot{z} = eEz$$

From the third equation, it is noted that the charge moves with uniform acceleration in the z-direction and is given by

$$Z = \frac{eE_z}{2m} t^2 + v_{0z}t \quad \dots\dots\dots 1.13$$

Multiplying the 2nd equation by i and combining with the first, we get

$$\frac{d}{dt}(\dot{x} + i\dot{y}) + i\omega(\dot{x} + i\dot{y}) = i \frac{e}{m} E_y$$

or
$$\dot{x} + i\dot{y} = \alpha e^{-i\omega t} + \frac{cE_y}{H} \quad \dots\dots\dots 1.14$$

Separating the real and imaginary parts we get

$$\dot{x} = a \cos \omega t + \frac{cE_y}{H} \quad \dots\dots\dots 1.15$$

$$\dot{y} = -\alpha \sin \omega t$$

The average velocity of the particle along x-axis and y-axis are

$$\bar{\dot{x}} = \frac{cE_y}{H}, \bar{\dot{y}} = 0 \quad \dots\dots\dots 1.16$$

Integrating and choosing the constant of integration so that at t= 0, x = y = 0 and we get

$$x = \frac{\alpha}{\omega} \sin \omega t + \frac{cE_y}{H} t \quad \dots\dots\dots 1.17$$

$$y = \frac{\alpha}{\omega} (\cos \omega t - 1)$$

These equations define a **trochoid**. Depending on whether a is large or smaller in absolute value than the quantity $\langle \dot{z} \rangle \approx v_0 + \omega_H(x) \approx \frac{v_0^2}{\omega_H R}$ the projection of the trajectory on the plane xy.

If
$$\alpha = \frac{-cE_y}{H}, \text{ then}$$

$$x = \frac{cE_y}{\omega H} (\omega t - \sin \omega t) \quad \dots\dots\dots 1.18$$

$$y = \frac{cE_y}{\omega H} (1 - \cos \omega t) \quad \dots\dots\dots 1.19$$

These gives the projection of the trajectory on the xy plane is a **cycloid**.

All the above formulas are valid for the velocity of the particle is small compared with the velocity of light and electric and magnetic fields satisfy the condition that

$$\frac{E_y}{H} \ll 1 \quad \dots\dots\dots 1.20$$

EXAMPLES:

I) Electric and Magnetic Field are Parallel:

To calculate the relativistic motion of a charged particle in parallel uniform electric and magnetic fields. In this case the magnetic field has

$$\dot{P}_x = \frac{e}{c} H v_y, \dot{P}_y = \frac{-e}{c} H v_x \quad \dots\dots\dots 1.21$$

$$z = \frac{\epsilon_0}{eE} \cosh \frac{E}{H} \phi \quad \dots\dots\dots 1.22$$

This gives the motion of the charged particle in **parametric form** and the trajectory is a helix with radius $\frac{cpt}{eH}$ and monotonically increasing step, along which the particle moves with decreasing angular velocity

$$\dot{\phi} = \frac{eHc}{\mathcal{E}_{kin}}$$

with a velocity along the z-axis which tends toward the value c.

no influence on the motion along the common direction of E and H (along the z-axis) and hence only the influence of electric field.

So the equation of motion in the xy-plane will be

II) Electric and Magnetic Field are Mutually Perpendicular:

For this, the equation of motion for the charged particle in which H is along z-direction and E along y-direction and E = H will be

$$\frac{dp_x}{dt} = \frac{e}{c} E v_y, \frac{dp_y}{dt} = eE \left(1 - \frac{v_y}{c} \right) \quad \dots\dots\dots 1.23$$

$$\frac{dp_z}{dt} = 0$$

which gives $P_z = \text{constant}$

It gives the motion of the particle in **parametric form** (i.e. parameter P_y) where the velocity increases most rapidly in the direction perpendicular to E and H along X-axis.

1.2 MOTION IN NON-UNIFORM, STATIC MAGNETIC FIELDS

Let us consider a non-uniform static magnetic field which varies slowly with distance in such a manner that the usual perturbation theory can be applied to get approximate solutions. For this the distance over which \bar{H} changes appreciably in magnitude or direction must be much greater than the gyration radius of the particle.

As an example consider a magnetic field which is independent of z. In the X-Y plane the lines of force are not parallel but slightly curved with a radius of curvature R that is large compared with the gyration radius a. Due to the symmetry of the problem it is advantageous to use cylindrical co-ordinate (ρ, ϕ, Z) with the origin at the centre of curvature.

The magnetic induction depends on the ratio $\frac{R}{\rho}$ and has only the ϕ component

$$H_\phi = H_0 \left(\frac{R}{\rho} \right) \quad \dots\dots\dots 1.24$$

The Lorentz force equation

$$m\ddot{\vec{r}} = e \left(\vec{E} + \vec{v} * \vec{H} \right) \quad \dots\dots\dots 1.25$$

becomes in cylindrical coordinates for the above magnetic field.

$$\ddot{\rho} - \rho \dot{\phi}^2 = -\omega_H \frac{R}{\rho} \dot{z} \quad \dots\dots\dots 1.26$$

$$\rho \ddot{\phi} + 2\dot{\rho}\dot{\phi} = 0 \quad \dots\dots\dots 1.27$$

$$\ddot{z} = \omega_H \ln\left(\frac{R}{\rho}\right) + \nu_0 \quad \dots\dots\dots 1.28$$

The equation (1.27) can be written as

$$\frac{d}{dt}(\rho^2 \dot{\phi}) = 0 \quad \dots\dots\dots 1.29$$

We obtain $\rho^2 \dot{\phi} = a$ constant which we write as $R\nu_0$

Had the magnetic field been a constant the trajectory would have been a helix, since H is not uniform, but does not change drastically we expect that ρ would be have a value close to R, when the trajectory of the helix has a radius small compared

to R. So ρ can be put as $\rho=R+x$ and $f\left(\frac{\rho}{R}\right)$ can

$$\ddot{x} + \left(\omega_H^2 + \frac{3}{R^2} \nu_0^2\right)x \approx \frac{\nu_0^2}{R} - \omega_H \nu_0 \quad \dots\dots\dots 1.30$$

This is the equation of motion of a harmonic oscillator around x with a displaced equilibrium position

$$\langle x \rangle \cong \frac{\nu_0^2}{\omega_H^2 R} - \frac{\nu_0}{\omega_H} \quad \dots\dots\dots 1.31$$

Here we have assumed $\nu_0 \ll \omega_H R$. The mean value of \dot{z} is

$$\langle \dot{z} \rangle \approx \nu_0 + \omega_H \langle x \rangle \approx \frac{\nu_0^2}{\omega_H R} \quad \dots\dots\dots 1.32$$

This is known as **Curvature drift**. If the spatial variation of the magnetic field is such that the gradient of the field is perpendicular to the direction of \vec{H} . then an analysis analogous to the above gives a gradient drift to velocity. Both then drifts are trouble some in confining high temperature plasmas, and the twisted figure eight toroidal design is made to keep the **plasma confined**.

II. CONCLUSION:

The above considerations are used in various ways such as cathode ray oscilloscopes and tubes, cyclotrons and other accelerators, motion of charged particles in the ionosphere, synchro-

cyclotron (Relativistic ion Cyclotron), $\frac{e}{m}$ of an

be expanded in powers of $\frac{x}{R}$, appropriately with

the approximation $\dot{z} \cong \omega_H x + \nu_0$ the radial equation of motion is approximately given by

electron by Thomson method, Thomson mass spectrograph, Aston's mass Spectrograph, Dempster, Mass Spectrograph, Magnetron Betatron, Hall effects etc.

BIBLIOGRAPHY:

- [1]. J. D. Jackson : Classical Electrodynamics, John Wiley (1999), Pergamon (1987)
- [2]. L. Landau and E. M. Lifshitz : Classical Theory of Fields.
- [3]. P. Achuthan, S Benjamin and K Venkatesan, J. Physics A 15, 3607 (1982).
- [4]. Brian G. Wybourne : Classical Groups for Physicists.
- [5]. K. Maharana, Math-Ph/0306089.
- [6]. A. P. Ruminez, J. Phys. - Condensed Matter.
- [7]. P. Drude Annalen der Physik 1, 566 and 3, 369 (1900)